

Entanglement Efficiencies in $\mathcal{PT}$ -Symmetric Quantum Mechanics

Christian Zielinski^{1,*} and Qing-hai Wang^{2,†}

¹*Institut für Theoretische Physik, Universität Heidelberg,
Philosophenweg 16, 69120 Heidelberg, Germany*

²*Department of Physics, National University of Singapore,
117542, Singapore*

(Dated: May 16, 2012)

The degree of entanglement is determined for an arbitrary state of a broad class of $\mathcal{PT}$ -symmetric bipartite composite systems. Subsequently we quantify the rate with which entangled states are generated and show that this rate can be characterized by a small set of parameters. These relations allow one in principle to improve the ability of these systems to entangle states. It is also noticed that many relations resemble corresponding ones in conventional quantum mechanics.

PACS numbers: 03.67.Bg, 03.65.Ud, 03.65.Ta

Keywords: entanglement generation, entanglement efficiency, entanglement entropy, $\mathcal{PT}$ -symmetric quantum mechanics

I. INTRODUCTION

In conventional quantum mechanics, one demands that the Hamiltonian H generating the time evolution has a real spectrum and that the corresponding time evolution operator U is unitary. These conditions are fulfilled if the Hamiltonian is Hermitian, i.e. $H = H^\dagger$, which is usually considered an axiom of quantum mechanics. However, the condition of Hermiticity can be weakened. In the class of so-called $\mathcal{PT}$ -symmetric Hamiltonians, one can ensure real eigenvalues and a unitary time evolution even for explicitly non-Hermitian Hamiltonians [2, 3]. A thorough review of the foundations of $\mathcal{PT}$ -symmetric quantum mechanics can be found in [1].

In the following we will investigate entanglement phenomena in bipartite systems within this framework. An entangled state is a quantum state where two or more degrees of freedom are intertwined, so that they are not independent anymore. In this context a couple of historical discussions took place and gave deep insights into the nature of quantum mechanics, like the *Einstein-Podolsky-Rosen* paradox [8]. These states have a wide range of applications, for example in quantum information theory and quantum computing. The question of entanglement generation and entanglement efficiencies for conventional quantum mechanics was addressed earlier in [7].

Using $\mathcal{PT}$ -symmetric quantum mechanics to describe entanglement phenomena was also done in [13]. Especially for the case of bipartite systems, relations for the degree of entanglement of given states and the entanglement capability of certain systems could be found. Although we confirmed many of these results, we have some different and new findings. For a particular initial state, as well as for general states, we give relations between the efficiency of the system to generate entangled states

and the parameters of the Hamiltonian describing the dynamics of the system.

We will first introduce a measure for entanglement and a certain class of $\mathcal{PT}$ -symmetric Hamiltonians. We then quantify the degree of entanglement of an arbitrary and of a generalized *Einstein-Podolsky-Rosen* state. Subsequently, we are dealing with the question of entanglement generation of a particular $\mathcal{PT}$ -symmetric state and generalize for arbitrary states. The question we try to answer is how to characterize the rate of entanglement generation for this class of systems.

II. BIPARTITE SYSTEMS

A. A measure for entanglement

Let $\mathcal{H}_{i=1,\dots,N}$ denote a set of Hilbert spaces. We call a state of a composite system $\mathcal{H} = \mathcal{H}_1 \otimes \dots \otimes \mathcal{H}_N$ entangled, if there is no decomposition of the form $|\Psi\rangle = |\chi_1\rangle \otimes \dots \otimes |\chi_N\rangle$ with suitable $|\chi_i\rangle \in \mathcal{H}_i$. In the following we restrict ourselves to bipartite systems, i.e. $N = 2$.

A measure of entanglement are the entropies

$$E(\Psi) = -\text{tr}_1(\rho_1 \log_2 \rho_1) = -\text{tr}_2(\rho_2 \log_2 \rho_2), \quad (1)$$

where $\rho_1 = \text{tr}_2 \rho$ and $\rho_2 = \text{tr}_1 \rho$ are the reduced density matrices, $\rho = |\Psi\rangle\langle\Psi|$ is the density matrix itself and tr_i denotes the partial trace over the i^{th} subsystem [12]. Here $E(\Psi) \in [0, 1]$ and $E(\Psi) = 0$ if and only if $|\Psi\rangle \in \mathcal{H}$ is not entangled.

B. General entanglement content

Consider the Hamiltonian

$$H = \begin{pmatrix} re^{i\Theta} & s \\ s & re^{-i\Theta} \end{pmatrix}, \quad r, s, \Theta \in \mathbb{R}, \quad (2)$$

with $s^2 \geq r^2 \sin^2 \Theta$. Observe that H is in general not represented by a Hermitian matrix, $H^\dagger \neq H$. But it

*Electronic address: zielinski@thphys.uni-heidelberg.de

†Electronic address: phywq@nus.edu.sg

obeys $\mathcal{PT}$ -symmetry with

$$\mathcal{P} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad (3)$$

and $\mathcal{T}$ complex conjugation [4] (see also erratum [5]), i.e. $[H, \mathcal{PT}] = 0$.

Define $\sin \varphi \equiv r/s \cdot \sin \Theta$ with $\varphi \in [-\pi/2, \pi/2]$. The simultaneous eigenstates of H and $\mathcal{PT}$ are given by

$$\begin{aligned} |\phi_+\rangle &= \frac{1}{\sqrt{2 \cos \varphi}} \begin{pmatrix} e^{i\varphi/2} \\ e^{-i\varphi/2} \end{pmatrix}, \\ |\phi_-\rangle &= \frac{i}{\sqrt{2 \cos \varphi}} \begin{pmatrix} e^{-i\varphi/2} \\ -e^{i\varphi/2} \end{pmatrix}, \end{aligned} \quad (4)$$

with eigenvalues $E_{\pm} = r \cos \Theta \pm s \cos \varphi$. These eigenstates are orthonormal with respect to the positive $\mathcal{CPT}$ -inner product $\langle \Psi | \Phi \rangle_{\mathcal{CPT}} \equiv \langle \Psi |_{\mathcal{CPT}} \cdot | \Phi \rangle$, where $\langle \Psi |_{\mathcal{CPT}} \equiv (\mathcal{CPT} | \Psi \rangle)^{\dagger}$, see [4]. For this Hamiltonian, the $\mathcal{C}$ operator has the form

$$\mathcal{C} = \begin{pmatrix} i \tan \varphi & \sec \varphi \\ \sec \varphi & -i \tan \varphi \end{pmatrix} \quad (5)$$

with $[H, \mathcal{C}] = 0$.

In the following we will consider the composite system $\mathcal{H} = \mathcal{H}_1 \times \mathcal{H}_2$ of Hilbert spaces $\mathcal{H}_{i=1,2}$ with dynamics governed by the Hamiltonian $H = H_1 \otimes H_2$, where H_i is of the form of (2) with $r_i, s_i, \Theta_i \in \mathbb{R}$, $s_i^2 \geq r_i^2 \sin^2 \Theta_i$, $\sin \varphi_i \equiv r_i/s_i \cdot \sin \Theta_i$ and $|\phi_{\pm}\rangle_i \in \mathcal{H}_i$ being the eigenstates of H_i for $i = 1, 2$. Our respective $\mathcal{C}$ operator reads $\mathcal{C}_1 \otimes \mathcal{C}_2$, where $\mathcal{C}_i$ is of the form of (5) with corresponding parameters. As

$$\begin{aligned} [H_1 \otimes H_2, \mathcal{PT} \otimes \mathcal{PT}] &= [H_1 \otimes H_2, \mathcal{C}_1 \otimes \mathcal{C}_2] \\ &= [\mathcal{C}_1 \otimes \mathcal{C}_2, \mathcal{PT} \otimes \mathcal{PT}] \\ &= 0 \end{aligned} \quad (6)$$

holds, H obeys $\mathcal{PT}$ symmetry and we can define a $\mathcal{CPT}$ -inner product as above. The decomposition of the operators means the same direct-product form of the metric operator η in [14]. According to proposition 2 and proposition 3 in the same paper, this permits a proper quantum mechanical description of the bipartite system with a unitary time evolution.

Now consider the density matrix $\rho = |\Psi(t)\rangle\langle\Psi(t)|_{\mathcal{CPT}}$ of a system in the state

$$\begin{aligned} |\Psi(t)\rangle &= \alpha(t)|\phi_+\rangle_1 \otimes |\phi_+\rangle_2 + \beta(t)|\phi_+\rangle_1 \otimes |\phi_-\rangle_2 \\ &\quad + \gamma(t)|\phi_-\rangle_1 \otimes |\phi_+\rangle_2 + \delta(t)|\phi_-\rangle_1 \otimes |\phi_-\rangle_2 \end{aligned} \quad (7)$$

with $|\Psi(t)\rangle \in \mathcal{H}$. The eigenvalues of $\rho_1(t) = \text{tr}_2 \rho(t)$ are given by $\lambda_{\pm}(t) = \frac{1}{2} \pm \frac{1}{2} \sqrt{1 - \Xi(t)}$ with $\Xi(t) = 4|\alpha(t)\delta(t) - \beta(t)\gamma(t)|^2$, which is a simplification of the result in [13]. Hence the entanglement content is

$$\begin{aligned} E(t) &\equiv E(\Psi(t)) \\ &= -\lambda_+(t) \log_2 \lambda_+(t) - \lambda_-(t) \log_2 \lambda_-(t). \end{aligned} \quad (8)$$

Note that $|\Psi(t)\rangle$ only separates if $\alpha(t_0)\delta(t_0) = \beta(t_0)\gamma(t_0)$ for $t = t_0$.




Figure 1: Entanglement content $E(\Psi^-)$ as a function of φ_1 and φ_2 .

C. The Einstein-Podolsky-Rosen state

Consider the *Einstein-Podolsky-Rosen* state from conventional quantum mechanics and normalize them with respect to the $\mathcal{CPT}$ -inner product:

$$|\uparrow\rangle_i = \sqrt{\cos \varphi_i} \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad |\downarrow\rangle_i = \sqrt{\cos \varphi_i} \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \quad (9)$$

Overall normalization yields

$$|\Psi^-\rangle = \kappa (|\uparrow\rangle_1 \otimes |\downarrow\rangle_2 - |\downarrow\rangle_1 \otimes |\uparrow\rangle_2) \quad (10)$$

with $\kappa = [2(1 - \sin \varphi_1 \sin \varphi_2)]^{-1/2}$. The eigenvalues of the reduced density matrix of the first subsystem are given by

$$\lambda_{\pm} = \frac{1}{2} \pm \frac{\sin \varphi_1 - \sin \varphi_2}{2(1 - \sin \varphi_1 \sin \varphi_2)}. \quad (11)$$

Our results are in disagreement with [13] mainly due to the author's use of non- $\mathcal{CPT}$ -normalized states.

We can consider the entanglement content

$$E(\Psi^-) = -\lambda_+ \log_2 \lambda_+ - \lambda_- \log_2 \lambda_- \quad (12)$$

as a function of φ_1 and φ_2 , i.e. of the Hamiltonians H_1 and H_2 . The result can be seen in figure 1. Note the cases $\varphi_1 = \varphi_2$ or $\varphi_1 + \varphi_2 = \pm\pi$, where (10) has the form of a $\mathcal{PT}$ -symmetric *Bell* state.




Figure 2: Entanglement content $E(t)$ (solid) and entanglement rate $\Gamma(t)/\tau^{-1}$ (dashed) of $|\Psi(t)\rangle$ with $\tau = 2\hbar/(\omega_1\omega_2)$.

III. ENTANGLEMENT GENERATION

A. Entanglement capability

We investigate the question how to increase the capability of a system to entangle states. More precisely we want to understand the dependencies of the entanglement capability $\Gamma(t) \equiv dE(t)/dt$ of the parameters of the system. In order to determine the time evolution operator of the Hamiltonian $H_1 \otimes H_2$ (see section II B) define

$$\vec{n}_i \equiv \frac{2}{\omega_i} (s_i, 0, i r_i \sin \Theta_i)^\top, \quad \omega_i \equiv 2s_i \cos \varphi_i. \quad (13)$$

Rewrite H as

$$H = \left(r_1 \cos \Theta_1 \mathbb{1} + \frac{\omega_1}{2} \vec{n}_1 \cdot \vec{\sigma} \right) \otimes \left(r_2 \cos \Theta_2 \mathbb{1} + \frac{\omega_2}{2} \vec{n}_2 \cdot \vec{\sigma} \right), \quad (14)$$

with $\vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)^\top$, σ_i denoting the Pauli matrices and $\mathbb{1}$ the 2×2 identity matrix. Expanding the expression for H yields four terms of which only one can generate entanglement. We restrict ourselves to this term and define $\hat{H} = \omega_1\omega_2/4 \cdot (\vec{n}_1 \cdot \vec{\sigma} \otimes \vec{n}_2 \cdot \vec{\sigma})$ resulting in a time evolution operator $U(t) = \exp(-i\hat{H}t/\hbar)$ given by

$$U(t) = \cos\left(\frac{\omega_1\omega_2 t}{4\hbar}\right) (\mathbb{1} \otimes \mathbb{1}) - i \sin\left(\frac{\omega_1\omega_2 t}{4\hbar}\right) (\vec{n}_1 \cdot \vec{\sigma} \otimes \vec{n}_2 \cdot \vec{\sigma}). \quad (15)$$

Consider now a simple initial state, namely $|\Psi(t=0)\rangle = |\uparrow\rangle \otimes |\uparrow\rangle$, and apply $U(t)$ to find

$$|\Psi(t)\rangle = \alpha(t) |\uparrow\rangle \otimes |\uparrow\rangle + \beta(t) |\uparrow\rangle \otimes |\downarrow\rangle + \gamma(t) |\downarrow\rangle \otimes |\uparrow\rangle + \delta(t) |\downarrow\rangle \otimes |\downarrow\rangle \quad (16)$$

with

$$\alpha(t) = \cos\left(\frac{\omega_1\omega_2 t}{4\hbar}\right) + \frac{4ir_1r_2 \sin \Theta_1 \sin \Theta_2}{\omega_1\omega_2} \sin\left(\frac{\omega_1\omega_2 t}{4\hbar}\right),$$

$$\begin{aligned} \beta(t) &= \frac{4s_2r_1 \sin \Theta_1}{\omega_1\omega_2} \sin\left(\frac{\omega_1\omega_2 t}{4\hbar}\right), \\ \gamma(t) &= \frac{4s_1r_2 \sin \Theta_2}{\omega_1\omega_2} \sin\left(\frac{\omega_1\omega_2 t}{4\hbar}\right), \\ \delta(t) &= -\frac{4is_1s_2}{\omega_1\omega_2} \sin\left(\frac{\omega_1\omega_2 t}{4\hbar}\right). \end{aligned} \quad (17)$$

Up to here our results are in agreement with [13]. We now find the entanglement content to be given by

$$E(t) = -\lambda_+(t) \log_2 \lambda_+(t) - \lambda_-(t) \log_2 \lambda_-(t) \quad (18)$$

with

$$\lambda_{\pm}(t) = \frac{1}{2} \pm \frac{1}{2} \cos\left(\frac{\omega_1\omega_2 t}{2\hbar}\right). \quad (19)$$

We find the entanglement rate to be

$$\begin{aligned} \Gamma(t) &= \frac{dE(t)}{dt} \\ &= \frac{\omega_1\omega_2}{\hbar} \sin\left(\frac{\omega_1\omega_2 t}{2\hbar}\right) \log_{16} \cot^2\left(\frac{\omega_1\omega_2 t}{4\hbar}\right). \end{aligned} \quad (20)$$

The maximal entanglement capability is $\Gamma_{\max} = \max_t \Gamma(t) = 0.4781 \omega_1\omega_2/\hbar$. Therefore by changing ω_i due to an adjustment of the parameters of the Hamiltonians H_i , we can control $\Gamma_{\max}$. The typical time dependency of the entanglement content and entanglement rate can be seen in figure 2.

B. Efficiency of general systems

If we have a given Hamiltonian one may ask how to maximize the entanglement rate of the system. We want to generalize some results of conventional quantum mechanics addressed in [7] to the $\mathcal{PT}$ -symmetric case.

Consider an arbitrary $\mathcal{PT}$ -symmetric two qubit system. Using the *Schmidt*-decomposition theorem we rewrite an arbitrary state $|\Psi(t)\rangle \in \mathcal{H}_1 \times \mathcal{H}_2$ as

$$|\Psi(t)\rangle = \sqrt{p(t)} |\varphi_t\rangle \otimes |\chi_t\rangle + e^{i\alpha} \sqrt{1-p(t)} |\varphi_t^\perp\rangle \otimes |\chi_t^\perp\rangle \quad (21)$$

with $p \in [0, 1]$ as *Schmidt*-coefficient and $\langle \varphi_t | \varphi_t^\perp \rangle_{\mathcal{CP}\mathcal{T}} = \langle \chi_t | \chi_t^\perp \rangle_{\mathcal{CP}\mathcal{T}} = 0$ as *Schmidt*-vectors. The entanglement content is given by

$$E(t) = -p(t) \log_2 p(t) - (1-p(t)) \log_2 (1-p(t)) \quad (22)$$

and the entanglement rate factorizes in two terms, i.e. $\Gamma(t) = dE(t)/dp(t) \times dp(t)/dt$, where

$$\frac{dE(t)}{dp(t)} = \frac{2}{\log 2} \operatorname{arctanh}(1-2p(t)). \quad (23)$$

After choosing the phase α appropriately the evolution of the *Schmidt*-coefficient is determined by the differential equation

$$\frac{dp(t)}{dt} = \frac{2}{\hbar} \sqrt{p(t)(1-p(t))} \times |(\langle \varphi_t |_{\mathcal{CPT}} \otimes \langle \chi_t |_{\mathcal{CPT}}) H(|\varphi_t^\perp\rangle \otimes |\chi_t^\perp\rangle)|. \quad (24)$$

This relation is known from conventional quantum mechanics, but also holds for $\mathcal{PT}$ -symmetric systems. Define

$$\hbar\Omega \equiv \max_{\substack{\|\varphi\|=1 \\ \|\chi\|=1}} |(\langle \varphi |_{\mathcal{CPT}} \otimes \langle \chi |_{\mathcal{CPT}}) H(|\varphi^\perp\rangle \otimes |\chi^\perp\rangle)|. \quad (25)$$

Then the time evolution of $p(t)$ for an optimally prepared setup, i.e. a setup forcing the qubit states to be optimal at every instant of time

$$|(\langle \varphi_t |_{\mathcal{CPT}} \otimes \langle \chi_t |_{\mathcal{CPT}}) H(|\varphi_t^\perp\rangle \otimes |\chi_t^\perp\rangle)| = \hbar\Omega \quad (26)$$

for all t , follows from the differential equation

$$\frac{dp_{\text{opt}}(t)}{dt} = 2\Omega \sqrt{p_{\text{opt}}(t)(1-p_{\text{opt}}(t))}. \quad (27)$$

We find $p_{\text{opt}}(t) = \sin^2(\Omega t + \delta_0)$ with an integration constant δ_0 , which is in agreement with results from conventional quantum mechanics in [7] (see also [13]). Hence we can characterize the entanglement rate of an optimally prepared setup completely in terms of Ω via

$$\begin{aligned} \Gamma_{\text{opt}}(t) &= \frac{dp_{\text{opt}}(t)}{dt} \cdot \left. \frac{dE(t)}{dp(t)} \right|_{p_{\text{opt}}(t)} \\ &= \frac{2\Omega}{\log 2} \operatorname{arctanh}(\cos(2\Omega t + 2\delta_0)) \sin(2\Omega t + 2\delta_0). \end{aligned} \quad (28)$$

and find for the maximal rate $\Gamma_{\text{max}} = \max_t \Gamma_{\text{opt}}(t) = 1.9123\Omega$. The parameter Ω is the critical value one needs to maximize to efficiently entangle states. We remark that this result also holds in conventional quantum mechanics, where Ω is defined with conventional conjugates instead of $\mathcal{CPT}$ -conjugates.

IV. CONCLUSIONS

For the case of a state from a bipartite system we determined the degree of entanglement and saw the emergence of symmetrical patterns (see figure 1) in the case of the *Einstein-Podolsky-Rosen* state. We quantified the capability of a given $\mathcal{PT}$ -symmetric system to generate entangled states in terms of the parameters of the Hamiltonian. Their ability to entangle states can be described by the parameters $\omega_{i=1,2}$ in (13) or in general by Ω in (25). Many relations are similar to the corresponding ones in conventional quantum mechanics after replacing the usual inner product with the $\mathcal{CPT}$ -inner product. However, these results are not obvious and need to be checked. For example, the recent discussion of the quantum brachistochrone problem showed that $\mathcal{PT}$ -symmetric quantum mechanics can give some surprising results [6, 9–11].

Acknowledgments

This work was carried out at the Department of Physics of the National University of Singapore, whose hospitality is gratefully acknowledged. The results were obtained within the UROPS program and we thank the National University of Singapore, the University of Heidelberg and the German Academic Foundation for financial support.

-
- [1] CARL M BENDER, *Making sense of non-hermitian hamiltonians*, Reports on Progress in Physics, 70 (2007), p. 947.
 - [2] CARL M. BENDER AND STEFAN BOETTCHER, *Real spectra in non-hermitian hamiltonians having $\mathcal{PT}$ symmetry*, Phys. Rev. Lett., 80 (1998), pp. 5243–5246.
 - [3] CARL M. BENDER, STEFAN BOETTCHER, AND PETER N. MEISINGER, *$\mathcal{PT}$ -symmetric quantum mechanics*, Journal of Mathematical Physics, 40 (1999), pp. 2201–2229.
 - [4] CARL M. BENDER, DORJE C. BRODY, AND HUGH F. JONES, *Complex extension of quantum mechanics*, Phys. Rev. Lett., 89 (2002), p. 270401.
 - [5] ———, *Erratum: Complex extension of quantum mechanics [phys. rev. lett. 89, 270401 (2002)]*, Phys. Rev. Lett., 92 (2004), p. 119902.
 - [6] CARL M. BENDER, DORJE C. BRODY, HUGH F. JONES, AND BERNHARD K. MEISTER, *Faster than hermitian quantum mechanics*, Phys. Rev. Lett., 98 (2007), p. 040403.
 - [7] W. DÜR, G. VIDAL, J. I. CIRAC, N. LINDEN, AND S. POPESCU, *Entanglement capabilities of nonlocal hamiltonians*, Phys. Rev. Lett., 87 (2001), p. 137901.
 - [8] A. EINSTEIN, B. PODOLSKY, AND N. ROSEN, *Can quantum-mechanical description of physical reality be considered complete?*, Phys. Rev., 47 (1935), pp. 777–780.
 - [9] UWE GÜNTHER AND BORIS F. SAMSONOV, *$\mathcal{PT}$ -symmetric brachistochrone problem, lorentz boosts, and nonunitary operator equivalence classes*, Phys. Rev. A, 78 (2008), p. 042115.
 - [10] ———, *Naimark-dilated $\mathcal{PT}$ -symmetric brachistochrone*, Phys. Rev. Lett., 101 (2008), p. 230404.
 - [11] ALI MOSTAFAZADEH, *Hamiltonians generating optimal-speed evolutions*, Phys. Rev. A, 79 (2009), p. 014101.

- [12] M.A. NIELSEN AND I.L. CHUANG, *Quantum Computation and Quantum Information: 10th Anniversary Edition*, Quantum Computation and Quantum Information, Cambridge University Press, 2011.
- [13] ARUN PATI, *Entanglement in non-hermitian quantum theory*, Pramana, 73 (2009), pp. 485–498. 10.1007/s12043-009-0101-0.
- [14] G SCOLARICI AND L SOLOMBRINO, *Alternative descriptions and bipartite compound quantum systems*, Journal of Physics A: Mathematical and Theoretical, 42 (2009), p. 055303.